



## Anti-DKK1 monoclonal antibody, clone 252246 (DCABY-3999)

This product is for research use only and is not intended for diagnostic use.

## **PRODUCT INFORMATION**

Antigen Description	Members of the dickkopf-related protein family (Dkk-1, -2, -3, and -4) are secreted proteins with two cysteine-rich domains separated by a linker region. Dkk-3 and -4 also have one prokineticin domain. Dkk-1, -2, and -4 function as antagonists of canonical Wnt signaling by binding to LRP5/6, preventing interaction of LRP5/6 with Wnt-Frizzled complexes. Dkk-1, -2, and -4 also bind cell surface Kremen-1 or -2 and promote the internalization of LRP5/6. Antagonistic activity of Dkk-3 has not been demonstrated. Dkk proteins have distinct patterns of expression in adult and embryonic tissues and have a wide range of effects on tissue development and morphogenesis. The Dkk family also includes Soggy, which is homologous to Dkk-3 but not to the other family members. Soggy has not been shown to inhibit Wnt signaling, and its role in the pathway is unclear.
Specificity	Detects human Dkk-1 in ELISAs. Insandwich immunoassays, no cross-reactivity or interference with recombinant human (rh)Dkk-4, rhKremen-1, rhKremen-2 or rhLRP-6 is observed.
Immunogen	S. frugiperda insect ovarian cell line Sf 21-derived recombinant human Dkk-1. Thr32-His266 Accession Number O94907
Isotype	lgG2b
Source/Host	Mouse
Species Reactivity	Human
Clone	252246
Purification	Protein A or G purified from hybridoma culture supernatant
Conjugate	Unconjugated
Applications	Immunohistochemistry, ELISA Capture (Matched Pair)

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Email: info@creative-diagnostics.com

© Creative Diagnostics All Rights Reserved

Format	Liquid
Size	500 μg
Buffer	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose.
Preservative	None
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.  12 months from date of receipt, -20 to -70 °C as supplied.  1 month from date of receipt, 2 to 8 °C, reconstituted.  6 months from date of receipt, -20 to -70 °C, reconstituted.

## **GENE INFORMATION**

Gene Name	DKK1 dickkopf WNT signaling pathway inhibitor 1 [ Homo sapiens (human) ]
Official Symbol	DKK1
Synonyms	DKK1; dickkopf WNT signaling pathway inhibitor 1; SK; DKK-1; dickkopf-related protein 1; hDkk-1; dickkopf-1 like; dickkopf 1 homolog; dickkopf-like protein 1; dickkopf related protein-1;
Entrez Gene ID	<u>22943</u>
Protein Refseq	<u>NP_036374</u>
UniProt ID	<u>I1W660</u>
Chromosome Location	10q11.2
Pathway	Cardiac Progenitor Differentiation; Direct p53 effectors; Presenilin action in Notch and Wnt signaling; Regulation of Wnt-mediated beta catenin signaling and target gene transcription; Signal Transduction; Signaling by Wnt; TCF dependent signaling in resp
Function	growth factor activity; low-density lipoprotein particle receptor binding; protein binding; receptor antagonist activity; signal transducer activity;