



# Anti-PROK1 monoclonal antibody, clone 299709 (DCABY-4133)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Antigen Description</b>	Endocrine gland-derived vascular endothelial growth factor (EG-VEGF), also called prokineticin 1 (PK1), is a member of the prokineticin family of secreted proteins that share a common structural motif containing ten conserved cysteine residues that form five pairs of disulfide bonds.
<b>Specificity</b>	Detects human EG-VEGF/PK1 in direct ELISAs and Western blots.
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human EG-VEGF/PK1. Ala20-Phe105 Accession Number P58294
<b>Isotype</b>	IgG1
<b>Source/Host</b>	Mouse
<b>Species Reactivity</b>	Human
<b>Clone</b>	299709
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	ELISA Capture (Matched Pair)
<b>Format</b>	Liquid
<b>Size</b>	500 µg
<b>Buffer</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.
<b>Preservative</b>	None

<b>Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
	12 months from date of receipt, -20 to -70 °C as supplied.
	1 month, 2 to 8 °C under sterile conditions after reconstitution.
	6 months, -20 to -70 °C under sterile conditions after reconstitution.

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">PROK1 prokineticin 1 [ Homo sapiens (human) ]</a>
<b>Official Symbol</b>	PROK1
<b>Synonyms</b>	PROK1; prokineticin 1; PK1; PRK1; EGVEGF; prokineticin-1; EG-VEGF; mambakine; black mamba toxin-related protein; endocrine-gland-derived vascular endothelial growth factor;
<b>Entrez Gene ID</b>	<a href="#">84432</a>
<b>Protein Refseq</b>	<a href="#">NP_115790</a>
<b>UniProt ID</b>	<a href="#">A0A024R0B1</a>
<b>Chromosome Location</b>	1p21
<b>Pathway</b>	Class A/1 (Rhodopsin-like receptors); G alpha (q) signalling events; GPCR downstream signaling; GPCR ligand binding; Gastrin-CREB signalling pathway via PKC and MAPK; Peptide ligand-binding receptors; Signal Transduction; Signaling by GPCR;
<b>Function</b>	growth factor activity;