



Anti-FRZB monoclonal antibody, clone 244042 (DCABY-4138)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Antigen Description	Secreted Frizzled related proteins are secreted glycoproteins with homology to the Wnt-binding domain of the Frizzled family of transmembrane receptors. sFRPs contain an N-terminal Frizzled-like cysteine-rich domain (CRD) with 10 conserved cysteines and a C-terminal heparin-binding region. sFRPs function as soluble antagonists of Wnt signaling by binding directly to Wnt proteins and preventing their interactions with Frizzled receptors.
Specificity	Detects human and mousesFRP-3 in ELISAs and Western blots.
Immunogen	Mouse myeloma cell line NS0-derived recombinant human sFRP-3. Ala33-Asn325 Accession Number AAB51298
Isotype	IgG1
Source/Host	Mouse
Species Reactivity	Human
Clone	244042
Purification	Protein A or G purified from hybridoma culture supernatant
Conjugate	Unconjugated
Applications	Western Blot, ELISA Capture (Matched Pair)
Format	Liquid
Size	500 μg
Buffer	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose.

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Preservative	None
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
	12 months from date of receipt, -20 to -70 °C as supplied.
	1 month, 2 to 8 °C under sterile conditions after reconstitution.
	6 months, -20 to -70 °C under sterile conditions after reconstitution.

GENE INFORMATION

Gene Name	FRZB frizzled-related protein [Homo sapiens (human)]
Official Symbol	FRZB
Synonyms	FRZB; frizzled-related protein; FRE; OS1; FZRB; hFIZ; FRITZ; FRP-3; FRZB1; SFRP3; SRFP3; FRZB-1; FRZB-PEN; secreted frizzled-related protein 3; sFRP-3; frezzled; frizzled homolog-related; frizzled-related protein 1;
Entrez Gene ID	<u>2487</u>
Protein Refseq	<u>NP_001454</u>
UniProt ID	D9ZGF6
Chromosome Location	2q32.1
Pathway	Adipogenesis; Endochondral Ossification;
Function	PDZ domain binding; Wnt-activated receptor activity; Wnt-protein binding;