



## Anti-Ubiquitin monoclonal antibody, clone 94567 (DCABY-4229)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

Antigen Description	Ubiquitin is a 76 amino acid (aa) protein that is ubiquitously expressed in all eukaryotic organisms. Ubiquitin is highly conserved with 96% aa sequence identity shared between human and yeast Ubiquitin, and 100% aa sequence identity shared between human and mouse Ubiquitin. In mammals, four Ubiquitin genes encode for two Ubiquitin-ribosomal fusion proteins and two poly-Ubiquitin proteins. Cleavage of the Ubiquitin precursors by deubiquitinating enzymes gives rise to identical Ubiquitin monomers each with a predicted molecular weight of 8.6 kDa. Conjugation of Ubiquitin to target proteins involves the formation of an isopeptide bond between the C-terminal glycine residue of Ubiquitin and a lysine residue in the target protein. This process of conjugation, referred to as ubiquitination or ubiquitylation, is a multi-step process that requires three enzymes: a Ubiquitin-activating (E1) enzyme, a Ubiquitin-conjugating (E2) enzyme, and a Ubiquitin ligase (E3). Ubiquitination is classically recognized as a mechanism to target proteins for degradation and as a result, Ubiquitin was originally named ATP-dependent Proteolysis Factor 1 (APF-1). In addition to protein degradation, ubiquitination has been shown to mediate a variety of biological processes such as signal transduction, endocytosis, and post-endocytic sorting.
Specificity	Detects human Ubiquitin+1 in ELISAs and Western blots. In in ELISAs and Western blots, no cross-reactivity with Ubiquitin is observed.
Immunogen	E. coli-derived recombinant human Ubiquitin+1. Met1-Gly75, YADLREDPDRQDHHPGSGAQ (C-terminal sequence generated by a frameshift) Accession Number CAA44911
Isotype	lgG2b
Source/Host	Mouse
Species Reactivity	Human
Clone	94567

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Purification	Protein A or G purified from hybridoma culture supernatant
Conjugate	Unconjugated
Applications	Western Blot, ELISA Capture (Matched Pair)
Format	Liquid
Size	500 μg
Buffer	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose.
Preservative	None
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.  12 months from date of receipt, -20 to -70 °C as supplied.  1 month, 2 to 8 °C under sterile conditions after reconstitution.  6 months, -20 to -70 °C under sterile conditions after reconstitution.

## **GENE INFORMATION**

Synonyms  UBB; ubiquitin B; polyubiquitin B;  Entrez Gene ID  7314  Protein Refseq  NP 001268645  UniProt ID  P0CG47  Chromosome Location  17p12-p11.2  Pathway  APC/C-mediated degradation of cell cycle proteins; APC/C:Cdc20 mediated degradation of Cyclin B; APC/C:Cdc20 mediated degradation of mitotic proteins; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cd	Gene Name	UBB ubiquitin B [ Homo sapiens (human) ]
Entrez Gene ID 7314  Protein Refseq NP 001268645  UniProt ID P0CG47  Chromosome Location 17p12-p11.2  Pathway APC/C-mediated degradation of cell cycle proteins; APC/C:Cdc20 mediated degradation of Cyclin B; APC/C:Cdc20 mediated degradation of Securin; APC/C:Cdc20 mediated degradation of mitotic proteins; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cd	Official Symbol	UBB
Protein Refseq  NP 001268645  UniProt ID  P0CG47  Chromosome Location  17p12-p11.2  Pathway  APC/C-mediated degradation of cell cycle proteins; APC/C:Cdc20 mediated degradation of Cyclin B; APC/C:Cdc20 mediated degradation of Securin; APC/C:Cdc20 mediated degradation of mitotic proteins; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cd	Synonyms	UBB; ubiquitin B; polyubiquitin-B; polyubiquitin B;
UniProt ID  P0CG47  Chromosome Location  17p12-p11.2  Pathway  APC/C-mediated degradation of cell cycle proteins; APC/C:Cdc20 mediated degradation of Cyclin B; APC/C:Cdc20 mediated degradation of Securin; APC/C:Cdc20 mediated degradation of mitotic proteins; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cd	Entrez Gene ID	<u>7314</u>
Chromosome Location  17p12-p11.2  Pathway  APC/C-mediated degradation of cell cycle proteins; APC/C:Cdc20 mediated degradation of Cyclin B; APC/C:Cdc20 mediated degradation of Securin; APC/C:Cdc20 mediated degradation of mitotic proteins; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cd	Protein Refseq	NP 001268645
Pathway  APC/C-mediated degradation of cell cycle proteins; APC/C:Cdc20 mediated degradation of Cyclin B; APC/C:Cdc20 mediated degradation of Securin; APC/C:Cdc20 mediated degradation of mitotic proteins; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cd	UniProt ID	P0CG47
Cyclin B; APC/C:Cdc20 mediated degradation of Securin; APC/C:Cdc20 mediated degradation of mitotic proteins; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cd	Chromosome Location	17p12-p11.2
Function protein binding;	Pathway	Cyclin B; APC/C:Cdc20 mediated degradation of Securin; APC/C:Cdc20 mediated degradation
	Function	protein binding;