



Anti-T. vaginalis Monoclonal antibody, Clone C990M (DMAB4432)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	Trichomonas vaginalis
Target	T. vaginalis
Immunogen	Disrupted Trichomonas vaginalis
Isotype	IgG3
Source/Host	Mouse
Species Reactivity	T. vaginalis
Clone	C990M
Affinity Constant	Not determined
Purification	90% pure. Protein A chromatography
Conjugate	Unconjugated
Applications	Suitable for use in IFA and ELISA. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded. Recommended antibody pair for sandwich immunoassay: Capture Detection DMAB4431 Suggested pair for testing (Capture - Detection): DMAB4432 - DMAB4431
Format	Purified, Liquid
Concentration	Lot specific (OD280nm, E0.1% = 1.3)

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Email: info@creative-diagnostics.com

Size	1 mg
Buffer	0.01M PBS, pH 7.2
Preservative	0.1% Sodium Azide
Storage	Short term (up to 2 weeks) store at 2-8°C. Long term, aliquot and store at -20°C. Avoid multiple freeze/thaw cycles

BACKGROUND

IntroductionTrichomonasvaginalisis an anaerobic, flagellated protozoan, a form of microorganism.

Theparasitic microorganism is the causative agent of trichomoniasis, and is themost common pathogenic protozoan infection of humans in industrialized countries. Infection rates between men and women are the same with womenshowing symptoms while infections in men are usually asymptomatic. Transmission takes place directly because the trophozoite does not have acyst. The WHO has estimated that 180 million cases of infection are acquired annually worldwide. The estimates for North America alone are between 5 and 8 million new infections each year, with an estimated rate of asymptomatic cases as high as 50%. Usually treatment consists of metronidazole and tinidazole. Trichomonas vaginalis is a protozoa which causes inflammation of the vaginal canal. It is considered to be a sexually transmitted disease.

Keywords T vaginalis ; TV ; Trichomonas vaginalis; Eukarya; Metamonada; Parabasalia; Trichomonadida;

Trichomonas; T. vaginalis

Email: info@creative-diagnostics.com