



Anti-FADS1 polyclonal antibody (DPAB-DC1854)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Antigen Description	The protein encoded by this gene is a member of the fatty acid desaturase (FADS) gene family. Desaturase enzymes regulate unsaturation of fatty acids through the introduction of double bonds between defined carbons of the fatty acyl chain. FADS family members are considered fusion products composed of an N-terminal cytochrome b5-like domain and a C-terminal multiple membrane-spanning desaturase portion, both of which are characterized by conserved histidine motifs. This gene is clustered with family members FADS1 and FADS2 at 11q12-q13.1; this cluster is thought to have arisen evolutionarily from gene duplication based on its similar exon/intron organization.
----------------------------	---

Immunogen	A synthetic peptide corresponding to human FADS1. The sequence is C-QPSFEPTKNKELTDE
Source/Host	Goat
Species Reactivity	Human
Purification	Antigen affinity purification
Conjugate	Unconjugated
Applications	WB (Tissue lysate), ELISA,
Format	Liquid
Concentration	0.5 mg/mL
Size	100 µg
Buffer	In Tris saline, pH 7.3 (0.5% BSA, 0.02% sodium azide)
Preservative	0.02% Sodium Azide

Storage

Store at -20°C. Aliquot to avoid repeated freezing and thawing.

GENE INFORMATION

Gene Name	FADS1 fatty acid desaturase 1 [Homo sapiens (human)]
Official Symbol	FADS1
Synonyms	FADS1; fatty acid desaturase 1; D5D; TU12; FADS6; FADSD5; LLCDL1; delta-5 desaturase; delta(5) desaturase; delta-5 fatty acid desaturase; delta(5) fatty acid desaturase; linoleoyl-CoA desaturase (delta-6-desaturase)-like 1;
Entrez Gene ID	3992
Protein Refseq	NP_037534
UniProt ID	O60427
Chromosome Location	11q12.2-q13.1
Pathway	Biosynthesis of unsaturated fatty acids; Fatty acid metabolism; Fatty acid, triacylglycerol, and ketone body metabolism; Metabolism
Function	C-5 sterol desaturase activity; heme binding; iron ion binding; oxidoreductase activity