



Anti-PIV type 1 Polyclonal antibody (DPAB0132)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

| | |
|---------------------------|---|
| Specificity | Parainfluenza 1, all structural antigens. Recognizes Sendai virus by ELISA. Does not cross-react with Para types 2 and 3, Influenza A, Influenza B, Respiratory Syncytial Virus, HSV1, HSV2, Adenovirus, CMV, Measles, Mumps and Rubella by indirect IFA. Does |
| Target | PIV type 1 |
| Immunogen | Cantell strain |
| Source/Host | Goat |
| Species Reactivity | PIV |
| Purification | Purified IgG fraction covalently coupled to a highly purified preparation of horseradish peroxidase (RZ3). Care is taken to ensure adequate conjugation while preserving maximum enzyme activity. Free enzyme is removed. Estimated HRP:IgG substitution is 2-3 |
| Conjugate | Unconjugated |
| Applications | Suitable for use in immunocytochemistry and ELISA applications. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded. |
| Format | Purified, Liquid. |
| Concentration | 1-2mg/ml (OD280nm, E0.1% = 1.4) |
| Size | 1 ml |
| Buffer | PBS containing 10mg/ml BSA |
| Preservative | None |

Storage

Short term (up to 6 months) store at 2-8°C. Long term, aliquot and store at -20°C. Avoid multiple freeze/thaw cycles.

BACKGROUND

Introduction

Human parainfluenza viruses (HPIV) were first discovered in the late 1950s. HPIV is genetically and antigenically divided into types 1 to 4. HPIV 1 to HPIV 3 are major causes of lower respiratory infections in infants, young children, the immunocompromised, the chronically ill, and the elderly. Each subtype can cause somewhat unique clinical diseases in different hosts. HPIV are enveloped and of medium size (150 to 250 nm), and their RNA genome is in the negative sense. These viruses belong to the Paramyxoviridae family, one of the largest and most rapidly growing groups of viruses causing significant human and veterinary disease. HPIV are closely related to recently discovered megamyxoviruses (Hendra and Nipah viruses) and metapneumovirus.

Keywords

HPIV 1; hPIV1; parainfluenza 1 virus; parainfluenza virus type; Parainfluenza; Parainfluenza Type 1; Parainfluenza 1; Mononegavirales; Paramyxoviridae
