



Anti-Gq α -Subunits (aa 115-133) polyclonal antibody (DPAB2914)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	Recognizes the Gq α -subunit in most tissues tested and recombinant Gq α -subunit expressed in Sf9 cells. Does not recognize the ~43 kDa G11 α -subunit in liver membrane or other members of the non-pertussis toxin-sensitive G-protein family.
Immunogen	a synthetic peptide (EVDVEKVSFAFENPYVDAIK) corresponding to amino acids 115-133 of the Gq α -subunit
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	N/A
Conjugate	Unconjugated
Applications	WB
Format	Liquid. In 140 mM NaCl, 100 mM potassium phosphate, pH 7.0.
Size	50 μ l
Preservative	None
Storage	$\leq -70^{\circ}\text{C}$, Avoid freeze/thaw

BACKGROUND

Introduction	Gq protein or Gq/11 is a heterotrimeric G protein subunit that activates phospholipase C (PLC). PLC in turn hydrolyzes Phosphatidylinositol 4,5-bisphosphate (PIP2) to diacyl glycerol (DAG)
---------------------	--

and inositol triphosphate (IP3) signal transduction pathway. DAG acts as a second messenger that activates Protein Kinase C (PKC) and IP3 helps in phosphorylation of some proteins.

Keywords

GNAQ; GAQ1; guanine nucleotide binding protein (G protein), q polypeptide; GAQ; G-ALPHA-q; guanine nucleotide-binding protein G(q) subunit alpha; guanine nucleotide-binding protein alpha-q; OTTHUMP00000021516
