



Anti-MMP10 polyclonal antibody [Biotin] (DPABY-451)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Antigen Description	The matrix metalloproteinases (MMPs) consist of 24 known human zinc proteases with essential roles in breaking down components of the extracellular matrix (ECM). Additional MMP substrates include cytokines, chemokines, growth factors and binding proteins, cell/cell adhesion molecules, and other proteinases. With a few exceptions, MMPs share common structural motifs including a pro-peptide domain, a catalytic domain, a hinge region, and a hemopexin-like domain. Synthesized as pro-enzymes, most MMPs are secreted before conversion to their active form. MMP activities are modulated on several levels including transcription, pro-enzyme activation, or by their endogenous inhibitors, tissue inhibitors of metalloproteinases (TIMPs). A subset of MMPs are associated with membranes and designated as membrane-type metalloproteinases (MT-MMP).
Specificity	Detects human MMP-10 in ELISAs and Western blots. In sandwich immunoassays,less than 0.1% cross-reactivity with recombinant human (rh) MMP-1, rhMMP-2, rhMMP-3, rhMMP-1, rhMMP-14, rhMMP-14, rhMMP-16, rhTIMP-1, rhTIMP-2, rhTIMP-3, and rhTIMP-4 is observed.
Immunogen	Mouse myeloma cell line NS0-derived recombinant human MMP-10 . Tyr18-Cys476 Accession Number P09238
Isotype	IgG
Source/Host	Goat
Species Reactivity	Human
Purification	Antigen Affinity-purified
Conjugate	Biotin
Applications	Western Blot, ELISA Detection (Matched Pair)

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Format	Liquid
Size	50 μg
Buffer	Lyophilized from a 0.2 μm filtered solution in PBS with BSA as a carrier protein.
Preservative	None
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

GENE INFORMATION

Gene Name	MMP10 matrix metallopeptidase 10 (stromelysin 2) [Homo sapiens (human)]
Official Symbol	MMP10
Synonyms	MMP10; matrix metallopeptidase 10 (stromelysin 2); SL-2; STMY2; stromelysin-2; MMP-10; transin 2; transin-2; stromelysin 2; matrix metalloprotease 10; matrix metalloproteinase-10; matrix metalloproteinase 10 (stromelysin 2);
Entrez Gene ID	4319
Protein Refseq	NP 002416
UniProt ID	<u>P09238</u>
Chromosome Location	11q22.3
Pathway	Activation of Matrix Metalloproteinases; Collagen degradation; Degradation of the extracellular matrix; Extracellular matrix organization; Matrix Metalloproteinases;
Function	calcium ion binding; metalloendopeptidase activity; zinc ion binding;

 ${\it Email: info@creative-diagnostics.com}$

Tel: 1-631-624-4882 Fax: 1-631-938-8221