



Anti-TAF2 (C-terminal) polyclonal antibody (DPATB-H82777)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	Rabbit Anti-TAF2 Polyclonal Antibody
Target	TAF2
Immunogen	Synthetic peptide corresponding to C terminal residues of Bakers Yeast (<i>Saccharomyces cerevisiae</i>) TAF2.
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	<i>Saccharomyces cerevisiae</i>
Purification	Immunogen affinity purified
Conjugate	Unconjugated
Applications	ELISA, WB
Cellular Localization	Nuclear
Format	Liquid
Size	50 µg
Buffer	50% Glycerol, PBS
Preservative	0.01% Sodium Azide
Storage	Store at +4°C short term (1-2 weeks). Aliquot and store at -20°C long term. Avoid repeated freeze / thaw cycles.

BACKGROUND

Introduction

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly,

Keywords

150 kDa cofactor of initiator function,150kDa,4732460C16Rik,AI425886,CIF150,Cofactor of initiator function, 150kD subunit,MGC166014,RNA polymerase II TBP associated factor subunit B 3,RNA polymerase II TBP associated factor subunit B,TAF(II)150,TAF2 RNA p

GENE INFORMATION

Entrez Gene ID

[6873](#)

UniProt ID

[B3KMD8](#)