



# Anti-Buprenorohine/NorBuprenorohine polyclonal antibody (DPBT-68302SB)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Product Overview</b>	Sheep Anti Buprenorphine/NorbuprenorphineSheep Anti Buprenorphine/Norbuprenorphine	
Specificity	DPBT-68302SB recognises Buprenorphine and Norbuprenorphine. Buprenorphine is a semi-synthetic opiate first marketed as an analgesic. It is prescribed for the treatment of moderate to severe pain or for anaesthetic induction. It is available in transdermal patches for long-term treatment of pain not responding to non-opioids. It is also used in opioid dependence treatment as withdrawal symptoms are milder than those of methadone. Norbuprenorphine is the active, dealkylated metabolite of buprenorphine. It acts as a stronger partial agonist against the mu opioid receptor than buprenorphine, but it does not cross the blood-brain barrier easily.	
Immunogen	Norbuprenorphine-bovine thyroglobulin.	
Isotype	IgG	
Source/Host	Sheep	
Species Reactivity	Chemical	
Conjugate	Unconjugated	
Applications	ELISA	
Preparation	Purified IgG prepared by caprylic acid and ammonium sulphate precipitation	
Format	Purified IgG - liquid	
Concentration	IgG concentration 1mg/ml	
Size	250 μg	
Buffer	Phosphate buffered saline	

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Email: info@creative-diagnostics.com

© Creative Diagnostics All Rights Reserved

Preservative	0.09% Sodium Azide

### Storage Storage in frost-free freezers is not recommended. This

product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation

before use.

## **BACKGROUND**

### Introduction Buprenorphine (C29 H41 NO4) is a prescription medication for people addicted to heroin or

other opiates that acts by relieving the symptoms of opiate withdrawal such as agitation, nausea and insomnia. Buprenorphine is more weakly addictive and has a lower risk of

overdose than methadone. The effects last for about three days.

## **Keywords** Buprenorphine; BUPRENORPHINE BASE; Buprenorphine 6,14-ethenomorphinan-7-

methanol,17-(cyclopropylmethyl)-alpha-(1,1-dimethylethy; I)-4,5-epoxy-18,19-dihydro-3-

hydroxy-6-methoxy-alpha-methyl-,(5-alpha,7-alpha; rx6029m;

BuprenorphineBase(Narcotic)C29H41No4,Edmf; Buprenorphine Base (Narcotic); Buprenex;

CL-112302; Lepetan; NIH-8805; Subutex; Temgesi; UM-952; BUPRENOPHINE;

Buprenorphine (base and/or unspecified salts); Methanol (test Buprenorphine,100ug/mL)