



Anti-HP polyclonal antibody (CABT-BL1827)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

| Immunogen | Affinity purified human haptoglobin. |
|-----------------------|---|
| Isotype | IgG |
| Source/Host | Goat |
| Species Reactivity | Human |
| Purification | Whole antiserum |
| Conjugate | Unconjugated |
| Applications | AP |
| Cellular Localization | Secreted. |
| Format | Liquid |
| Buffer | Tris buffered saline |
| Preservative | 0.1% Sodium Azide |
| Storage | Add glycerol to a final volume of 50%, aliquot and store at -20°C. Avoid repeated freeze / thaw cycles. |

BACKGROUND

Introduction

As a result of hemolysis, hemoglobin is found to accumulate in the kidney and is secreted in the urine. Haptoglobin captures, and combines with free plasma hemoglobin to allow hepatic recycling of heme iron and to prevent kidney damage. Haptoglobin also acts as an Antimicrobial; Antioxidant, has antibacterial activity and plays a role in modulating many aspects of the acute phase response. Hemoglobin/haptoglobin complexes are rapidely cleared

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

© Creative Diagnostics All Rights Reserved

by the macrophage CD163 scavenger receptor expressed on the surface of liver Kupfer cells through an endocytic lysosomal degradation pathway. Uncleaved haptoglogin, also known as zonulin, plays a role in intestinal permeability, allowing intercellular tight junction disassembly, and controlling the equilibrium between tolerance and immunity to non-self antigens.

GENE INFORMATION

| Entrez Gene ID | 3240 |
|----------------|---------------|
| Protein Refseq | NP_001119574 |
| UniProt ID | <u>P00738</u> |