



Anti-MLLT10 (C-terminal) polyclonal antibody (CABT-BL2444)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Immunogen	A synthetic peptide (conjugated with KLH) corresponding to C-terminus of human MLLT10.
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Purification	Protein G purification
Conjugate	Unconjugated
Applications	WB, IHC, ELISA
Format	Liquid
Buffer	In PBS (0.09% sodium azide)
Preservative	0.09% Sodium Azide
Storage	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.

BACKGROUND

Introduction

This gene encodes a transcriptional coactivator that plays an essential role in regulating gene expression during early development and hematopoiesis. The encoded protein contains multiple conserved functional domains. One of these domains, the SET domain, is responsible for its histone H3 lysine 4 (H3K4) methyltransferase activity which mediates chromatin

45-1 Ramsey Road, Shirley, NY 11967, USA

Email:info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

modifications associated with epigenetic transcriptional activation. This protein is processed by the enzyme Taspase 1 into two fragments, MLL-C and MLL-N. These fragments reassociate and further assemble into different multiprotein complexes that regulate the transcription of specific target genes, including many of the HOX genes. Multiple chromosomal translocations involving this gene are the cause of certain acute lymphoid leukemias and acute myeloid leukemias. Alternate splicing results in multiple transcript variants.

GENE INFORMATION

Entrez Gene ID	8028
Protein Refseq	NP_001182555
UniProt ID	<u>P55197</u>