



# Mouse Anti-Human CD32 (FcγRIIA) Monoclonal antibody, clone IV.3 (CABT-L4325)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

### Product Overview

The IV.3 monoclonal antibody reacts with human CD32 also known as Fc $\gamma$ RII and FCRII, a 40 kDa polymorphic transmembrane glycoprotein and an Ig superfamily member. CD32 is expressed on monocytes/macrophages, granulocytes, platelets and B cells. CD32 enables interaction between Fc  $\gamma$  RII-expressing cells and opsonized antigen or IgG-containing immune complexes. This allows CD32 to function in the activation or inhibition of immune responses including degranulation, phagocytosis, ADCC, cytokine release, and B cell proliferation. The IV.3 antibody has been shown to block the biological effects of CD32 in vitro. Additionally, IV.3 f(ab')2 fragments have been used to block CD32 in vivo in transgenic mice expressing human CD32.

<b>Target</b>	Human CD32 (Fc $\gamma$ RIIA)
<b>Immunogen</b>	Human K562 leukemia cell line
<b>Isotype</b>	IgG2b
<b>Source/Host</b>	Mouse
<b>Species Reactivity</b>	Human
<b>Clone</b>	IV.3
<b>Purification</b>	Protein G purified. Purity>95%. Determined by SDS-PAGE
<b>Conjugate</b>	Functional Grade
<b>Applications</b>	in vivo Fc $\gamma$ RIIA blockade in humanized mice, in vitro Fc $\gamma$ RIIA blockade, ELISA, FC
<b>Molecular Weight</b>	150 kDa

<b>Format</b>	0.2 µM filtered liquid. Purified from tissue culture supernatant in an animal free facility
<b>Concentration</b>	Lot specific
<b>Size</b>	5 mg
<b>Buffer</b>	PBS, pH 7.0. Contains no stabilizers or preservatives. [low endotoxin azide-free]  Endotoxin level: <2EU/mg (<0.002EU/µg). Determined by LAL gel clotting assay Related dilution buffer: CABT-LB04
<b>Preservative</b>	None
<b>Storage</b>	The antibody solution should be stored undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.
<b>Ship</b>	Wet ice

## BACKGROUND

<b>Introduction</b>	Binds to the Fc region of immunoglobulins gamma. Low affinity receptor. By binding to IgG it initiates cellular responses against pathogens and soluble antigens.
<b>Keywords</b>	ARH12;ARHA;H12;ras homolog gene family member A;ras homolog gene family member B;ras homolog gene family member C;Rho cDNA clone 12;RHO12;RHOA;RHOA;rhob;rhoc;RHOH12;Small GTP binding protein RhoA;Transforming protein RhoA;

## GENE INFORMATION

<b>Official Symbol</b>	CD32
<b>Synonyms</b>	ARH12; ARHA; H12; ras homolog gene family member A; ras homolog gene family member B; ras homolog gene family member C; Rho cDNA clone 12; RHO12; RHOA; RHOA; rhob; rhoc; RHOH12; Small GTP binding protein RhoA; Transforming protein RhoA;
<b>References</b>	Walsh, T. G., et al. (2015). "SDF-1alpha is a novel autocrine activator of platelets operating through its receptor CXCR4." <i>Cell Signal</i> 27(1): 37-46. PubMed;