



# Rabbit Anti-Human SDC1 Monoclonal Antibody, clone CQ7146 (CABT-Z208R)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Immunogen</b>	Synthetic peptide corresponding to CD138 residues within aa200-300 of CD138 was used as an immunogen.
<b>Isotype</b>	IgG
<b>Source/Host</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Clone</b>	CQ7146
<b>Purification</b>	ProA affinity purified IgG.
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	IHC-P Recommended concentration: IHC-P: 1:100-1:200
<b>Molecular Weight</b>	32 kDa
<b>Cellular Localization</b>	Membrane
<b>Positive Control</b>	Plasmacytoma
<b>Format</b>	Liquid
<b>Concentration</b>	Lot specific
<b>Size</b>	100 µl

<b>Buffer</b>	PBS 59%, Sodium azide 0.01%, Glycerol 40%, BSA 0.05%.
<b>Preservative</b>	0.01% Sodium azide
<b>Storage</b>	Store at -20 °C. Avoid freeze/thaw cycles.
<b>Ship</b>	Wet ice

## BACKGROUND

**Introduction** CD138, also designated syndecan-1, is a member of the syndecan family of four transmembrane spanning proteins capable of binding to heparan sulfate and chondroitin sulfate molecules. The syndecans' main functions are to control cell growth and differentiation as well as to maintain cell adhesion and cell migration. Under normal conditions CD138 is predominantly expressed on mature plasma cells and early preB-cells, while other haematolymphoid cells are negative. Various types of epithelial cells are also CD138 positive. Squamous epithelial cells show strong membranous and some cytoplasmic staining, except for the mature superficial squamous epithelial cells which are unstained. Mature mesenchymal and neural tissues are not stained. Among haematolymphoid neoplasms, CD138 is expressed in practically all cases of plasma cell malignancies. Among nonhaematolymphoid neoplasms the expression of CD138 is found in various types of carcinoma. In the following, the large majority of cases is CD138 positive: skin squamous cell carcinoma and basal cell carcinoma, colorectal adenocarcinoma, cholangiocarcinoma, transitional cell carcinoma, endometrial adenocarcinoma, ductal and lobular breast carcinoma, hepatocellular carcinoma, renal cell carcinoma, and lung adenocarcinoma. CD138 is a very sensitive and specific marker for identification of plasma cells and plasma cell differentiation within haematolymphoid tissues in benign and neoplastic conditions.

**Keywords** SDC1; syndecan 1; SDC; CD138; SYND1; syndecan; syndecan-1; CD138 antigen; syndecan proteoglycan 1; heparan sulfate proteoglycan fibroblast growth factor receptor

## GENE INFORMATION

<b>Gene Name</b>	SDC1
<b>Entrez Gene ID</b>	<a href="#">6382</a>
<b>UniProt ID</b>	<a href="#">P18827</a>