



Anti-Rat IgG polyclonal antibody [Dylight® 800] (CPBT-66037GR)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

| Product Overview | Goat anti Rat IgG (Mouse Adsorbed) antibody recognizes rat IgG. Cross-reactivity with mouse IgG has been minimised by adsorption. Flow Cytometry Use 50ul of the suggested working dilution to label 1x106 cells in 100ul. |
|--------------------|---|
| Specificity | IgG |
| Immunogen | Rat IgG |
| Isotype | IgG |
| Source/Host | Goat |
| Species Reactivity | Rat |
| Conjugate | Dylight 800 |
| Applications | FC; IF; WB |
| Size | 100 μg |
| Preservative | 0.09% Sodium Azide |
| Storage | in frost free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. |

GENE INFORMATION

Gene Name | Ighg Immunoglobulin heavy chain (gamma polypeptide) [Rattus norvegicus (Norway rat)]

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

| Official Symbol | Ighg |
|---------------------|--|
| Synonyms | IGHG; Immunoglobulin heavy chain (gamma polypeptide); RGD1359539; IGG; |
| Entrez Gene ID | <u>299354</u> |
| UniProt ID | P20760 |
| Chromosome Location | 6q32 |
| Pathway | Classical antibody-mediated complement activation; Complement cascade; Creation of C4 and C2 activators; FCGR activation; Fcgamma receptor (FCGR) dependent phagocytosis; Immune System; Initial triggering of complement; Innate Immune System; Regulation of actin dynamics for phagocytic cup formation; Role of phospholipids in phagocytosis; |
| Function | antigen binding |
| | |