



# Recombinant SARS-CoV-2 Spike RBD (A435S) [His] (DAGC190)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Product Overview</b>	A DNA sequence encoding the SARS-CoV-2 Spike Protein (RBD) (YP_009724390.1(A435S)) (Arg319-Phe541) was expressed with a polyhistidine tag at the C-terminus.
<b>Species</b>	SARS-CoV-2
<b>Purity</b>	> 95 % as determined by SDS-PAGE.
<b>Conjugate</b>	His
<b>Applications</b>	SDS-PAGE, ELISA
<b>Predicted N terminal</b>	Arg319
<b>Molecular Weight</b>	The recombinant SARS-CoV-2 Spike Protein (RBD, His Tag) consists of 234 amino acids and predicts a molecular mass of 26.56 kDa.
<b>Endotoxin</b>	< 1.0 EU per µg protein as determined by the LAL method.
<b>Format</b>	Lyophilized
<b>Size</b>	100 µg, 1 mg
<b>Buffer</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
<b>Preservative</b>	None
<b>Storage</b>	Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

# BACKGROUND

## Introduction

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process.

## Keywords

SARS-CoV-2; coronavirus; SARS-CoV-2 spike RBD; SARS-CoV-2 spike protein