



Anti-CAMLG monoclonal antibody (DCABH-10826)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Antigen Description	The immunosuppressant drug cyclosporin A blocks a calcium-dependent signal from the T-cell receptor (TCR) that normally leads to T-cell activation. When bound to cyclophilin B, cyclosporin A binds and inactivates the key signaling intermediate calcineurin. The protein encoded by this gene functions similarly to cyclosporin A, binding to cyclophilin B and acting downstream of the TCR and upstream of calcineurin by causing an influx of calcium. This integral membrane protein appears to be a new participant in the calcium signal transduction pathway, implicating cyclophilin B in calcium signaling, even in the absence of cyclosporin.
----------------------------	---

Immunogen	A synthetic peptide of human CAMLG is used for rabbit immunization.
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Purification	Protein A
Conjugate	Unconjugated
Applications	Western Blot (Transfected lysate); ELISA
Buffer	In 1x PBS, pH 7.4
Preservative	None
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

GENE INFORMATION

Gene Name	CAMLG calcium modulating ligand [Homo sapiens]
Official Symbol	CAMLG
Synonyms	CAMLG; calcium modulating ligand; calcium signal-modulating cyclophilin ligand; calcium modulating cyclophilin ligand; calcium signal modulating cyclophilin ligand; CAML; cyclophilin B binding protein; cyclophilin B-binding protein; calcium-modulating cyclophilin ligand; calcium-signal modulating cyclophilin ligand; MGC163197;
Entrez Gene ID	819
Protein Refseq	NP_001736
UniProt ID	P49069
Chromosome Location	5q23
Function	protein binding;