



## Anti-CYP2C8 monoclonal antibody (DCABH-11191)

This product is for research use only and is not intended for diagnostic use.

## **PRODUCT INFORMATION**

| Antigen Description | This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by phenobarbital. The enzyme is known to metabolize many xenobiotics, including the anticonvulsive drug mephenytoin, benzo(a)pyrene, 7-ethyoxycoumarin, and the anti-cancer drug taxol. This gene is located within a cluster of cytochrome P450 genes on chromosome 10q24. |
|---------------------|--|
| Immunogen           | A synthetic peptide of human CYP2C8 is used for rabbit immunization.   |
| Isotype             | IgG  |
| Source/Host         | Rabbit   |
| Species Reactivity  | Human  |
| Purification        | Protein A  |
| Conjugate           | Unconjugated   |
| Applications        | Western Blot (Transfected lysate); ELISA   |
| Buffer              | In 1x PBS, pH 7.4  |
| Preservative        | None   |
| Storage             | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.   |

## **GENE INFORMATION**

45-1 Ramsey Road, Shirley, NY 11967, USA

Email:info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

| Gene Name           | CYP2C8 cytochrome P450, family 2, subfamily C, polypeptide 8 [ Homo sapiens ]  |
|---------------------|--|
| Official Symbol     | CYP2C8   |
| Synonyms            | CYP2C8; cytochrome P450, family 2, subfamily C, polypeptide 8; cytochrome P450, subfamily IIC (mephenytoin 4 hydroxylase), polypeptide 8; cytochrome P450 2C8; CPC8; P450 form 1; cytochrome P450 IIC2; cytochrome P450 MP-12; cytochrome P450 MP-20; cytochrome P450 form 1; microsomal monooxygenase; xenobiotic monooxygenase; s-mephenytoin 4-hydroxylase; flavoprotein-linked monooxygenase; cytochrome P450, subfamily IIC (mephenytoin 4-hydroxylase), polypeptide 8; CYPIIC8; MP-12/MP-20; |
| Entrez Gene ID      | <u>1558</u>  |
| Protein Refseq      | NP 000761  |
| UniProt ID          | <u>P10632</u>  |
| Chromosome Location | 10q24.1  |
| Pathway             | Arachidonate Epoxygenase / Epoxide Hydrolase, organism-specific biosystem; Arachidonic acid metabolism, organism-specific biosystem; Arachidonic acid metabolism, conserved biosystem; Biological oxidations, organism-specific biosystem; Cytochrome P450 - arranged by substrate type, organism-specific biosystem; Drug metabolism - cytochrome P450, organism-specific biosystem; Drug metabolism - cytochrome P450, conserved biosystem;  |
| Function            | aromatase activity; caffeine oxidase activity; electron carrier activity; heme binding; metal ion binding; monooxygenase activity; oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen;  |