



Anti-PRKG1 monoclonal antibody, clone 0B5 (DCABH-845)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	Mouse monoclonal to cGKI
Antigen Description	<p>Serine/threonine protein kinase that acts as key mediator of the nitric oxide (NO)/cGMP signaling pathway. GMP binding activates PRKG1, which phosphorylates serines and threonines on many cellular proteins. Numerous protein targets for PRKG1 phosphorylation are implicated in modulating cellular calcium, but the contribution of each of these targets may vary substantially among cell types. Proteins that are phosphorylated by PRKG1 regulate platelet activation and adhesion, smooth muscle contraction, cardiac function, gene expression, feedback of the NO-signaling pathway, and other processes involved in several aspects of the CNS like axon guidance, hippocampal and cerebellar learning, circadian rhythm and nociception. Smooth muscle relaxation is mediated through lowering of intracellular free calcium, by desensitization of contractile proteins to calcium, and by decrease in the contractile state of smooth muscle or in platelet activation. Regulates intracellular calcium levels via several pathways: phosphorylates MRV11/IRAG and inhibits IP3-induced Ca(2+) release from intracellular stores, phosphorylation of KCNMA1 (BKCa) channels decreases intracellular Ca(2+) levels, which leads to increased opening of this channel. PRKG1 phosphorylates the canonical transient receptor potential channel (TRPC) family which inactivates the associated inward calcium current. Another mode of action of NO/cGMP/PKG1 signaling involves PKGI-mediated inactivation of the Ras homolog gene family member A (RhoA). Phosphorylation of RHOA by PRKG1 blocks the action of this protein in myriad processes: regulation of RHOA translocation; decreasing contraction; controlling vesicle trafficking, reduction of myosin light chain phosphorylation resulting in vasorelaxation. Activation of PRKG1 by NO signaling alters also gene expression in a number of tissues. In smooth muscle cells, increased cGMP and PRKG1 activity influence expression of smooth muscle-specific contractile proteins, levels of proteins in the NO/cGMP signaling pathway, down-regulation of the matrix proteins osteopontin and thrombospondin-1 to limit smooth muscle cell migration and phenotype. Regulates vasodilator-stimulated phosphoprotein (VASP) functions in platelets and smooth muscle.</p>
Immunogen	Recombinant full length Human cGKI produced in HEK293T cells (NP_006249).

Isotype	IgG2b
Source/Host	Mouse
Species Reactivity	Human
Clone	0B5
Purification	This antibody is purified from Mouse ascites fluid by affinity chromatography.
Conjugate	Unconjugated
Applications	WB, Flow Cyt, ICC/IF
Positive Control	HEK293T cell lysate transfected with pCMV6-ENTRY cGKI cDNA; COS7 cells transiently transfected by pCMV6-ENTRY cGKI; HEK293T cells transfected with cGKI overexpress plasmid; Jurkat cells.
Format	Liquid
Size	100 µl
Buffer	pH: 7.30; Preservative: 0.02% Sodium azide; Constituents: 48% PBS, 1% BSA, 50% Glycerol
Preservative	0.02% Sodium Azide
Storage	store at -20°C. Avoid repeated freeze / thaw cycles.
Ship	Shipped at 4°C.

GENE INFORMATION

Gene Name	PRKG1 protein kinase, cGMP-dependent, type I [Homo sapiens]
Official Symbol	PRKG1
Synonyms	PRKG1; protein kinase, cGMP-dependent, type I; PRKG1B, PRKGR1B; cGMP-dependent protein kinase 1; PGK; PKG; protein kinase, cGMP-dependent, regulatory, type I, beta; 1; cGK; cGK1; cGKI; cGK 1; PRKG1B; PRKGR1B; cGKI-BETA; cGKI-alpha; FLJ36117; MGC71944; DKF
Entrez Gene ID	5592
Protein Refseq	NP_001091982

UniProt ID	Q13976
Chromosome Location	10q11.2
Pathway	Adaptive Immune System, organism-specific biosystem; Gap junction, organism-specific biosystem; Gap junction, conserved biosystem; Hemostasis, organism-specific biosystem; Immune System, organism-specific biosystem; Long-term depression, organism-specific biosystem; Long-term depression, conserved biosystem;
Function	ATP binding; cGMP binding; cGMP-dependent protein kinase activity; calcium channel regulator activity; nucleotide binding; protein binding; protein serine/threonine kinase activity;