



Rabbit Anti-HBsAg PreS1 Polyclonal antibody (DMAB2924)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	Monoclonal Antibody to Adenovirus
Specificity	Specific for the hexon group antigen of many Adenovirus serotypes. Known reactivity with 34 serotypes of Adenovirus including types 40 and 41 (40, 41, 1, 1a, 2, 2c, 3, 3a, 4, 5, 5a, 5b, 5c, 5d, 6, 7, 7a, 8, 9, 10, 11, 12, 12a, 14, 16, 18, 19, 20, 26, 31, 34, 35, 36 and 37). Does not react with Influenza A, Influenza B, RSV, Parainfluenza 1, 2 & 3, Mycoplasma pneumoniae, H. pylori and Mammalian cells.
Target	Adenovirus
Immunogen	Infected cell extract with adenovirus type 6
Isotype	IgG1
Source/Host	Mouse
Species Reactivity	Adenovirus
Clone	A144
Affinity Constant	Not determined
Purification	>90% pure. Protein A chromatography
Conjugate	Unconjugated
Applications	Suitable for use in ELISA, Lateral flow and IFA. Not suitable for use in Western blot. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded. Recommended pairs for sandwich immunoassay:

- **Capture**
[DMAB2924](#)
[DMAB2924](#)
- **Detection**
[DPAB0138](#)
[DPAB0138](#)

Format	Purified, Liquid
Concentration	100ug/ml (OD280nm, E0.1% = 1.3)
Size	1 mg
Buffer	0.01M PBS, pH 7.2. This product contains no stabilizing proteins.
Preservative	0.1% Sodium Azide
Storage	Upon receipt, aliquot and store at -20°C. Avoid multiple freeze/thaw cycles.

BACKGROUND

Introduction Adenoviruses are DNA viruses generally widespread in nature that are frequently the cause of acute upper respiratory tract infections (i.e. common colds). Forty-seven known serotypes have been isolated since they were first discovered in 1953 with 3 types known to cause gastroenteritis. Several types have oncogenic potential though most cause self-limiting febrile illnesses characterised by inflammation of conjunctivae and the respiratory tract. The virus can be isolated from the majority of tonsils/adenoids surgically removed, indicating latent infections. It is not known how long the virus can persist in the body, or whether it is capable of reactivation after long periods. In patients experiencing immunosuppression (e.g. AIDS) it can be reactivated causing disease.

Keywords Adenovirus; Adenoviridae; Aviadenovirus; ADENOVIRUS; ADENOVIRUS F
