



# Anti-Adenovirus Hexon Monoclonal antibody, Clone C113M (DMAB2930)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Product Overview</b>	Monoclonal Antibody to Adenovirus hexon
Specificity	Specific for the hexon group antigen of Adenovirus. Reactive with Adenovirus types 1, 2, 3, 4, 5, 6, 7a, 8, 31, 40 and 41. Other types not tested. Does not react with Influenza A, Influenza B, RSV, Parainfluenza 1, 2 & 3, Mycoplasma pneumoniae, H. pylori and Mammalian cells
Target	Adenovirus Hexon
Immunogen	Adenovirus lysate
Isotype	lgG1
Source/Host	Mouse
Species Reactivity	Adenovirus
Clone	C113M
Affinity Constant	Not determined
Purification	>90% pure. Protein A chromatography.
Conjugate	Unconjugated
Applications	Suitable for use in ELISA and IFA. Not recommended for use with methanol-fixed cells. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded. Recommended pairs for sandwich immunoassay:  • Capture

45-1 Ramsey Road, Shirley, NY 11967, USA

Email:info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

DMAB2930

**DMAB2930** 

Detection

DMAB2931 DMAB2938

Suggested pair for testing (Capture - Detection): DMAB2930 - DMAB2938

Procedure	Matched Antibody Pairs
Format	Purified, Liquid
Concentration	100ug/ml (OD280nm, E0.1%= 1.3)
Size	1 mg
Buffer	0.01M PBS, pH 7.2. Product contains no stabilizing proteins.
Preservative	0.1% Sodium Azide
Storage	Upon receipt, store at -20°C. Avoid multiple freeze/thaw cycles.
Warnings	This product contains sodium azide, which has been classified as Xn (Harmful), in European Directive 67/548/EEC in the concentration range of 0.1–1.0%. When disposing of this reagent through lead or copper plumbing, flush with copious volumes of water to prevent azide build-up in drains.

### **BACKGROUND**

#### Introduction

Adenovirusesare DNA viruses generally widespread in nature that are frequently the cause of acute upper respiratory tract infections (i.e. common colds). Forty-sevenknown serotypes have been isolated since they were first discovered in 1953with 3 types known to cause gastroenteritis. Several types have oncogenic potential though most cause self-limiting febrile illnesses characterised by inflammation of conjunctivae and the respiratory tract. The virus can be isolated from the majority of tonsils/adenoids surgically removed, indicating latent infections. It is not known how long the virus can persist in the body, or whether it is capable of reactivation after long periods. In patients experiencing immunosuppression (e.g. AIDS) it can be reactivated causing disease.

#### Keywords

Adenovirus; Adenovirus Hexon; Adenovirus hexon; Hexon protein; Late protein 2; PII; Adenovirus; Adenoviridae; Aviadenovirus; ADENOVIRUS; ADENOVIRUS F

45-1 Ramsey Road, Shirley, NY 11967, USA

Email:info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221