

Anti-T. vaginalis Monoclonal antibody, Clone C986M (DMAB4429)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

| Specificity | Trichomonas vaginalis |
|--------------------|---|
| Target | T. vaginalis |
| Immunogen | Disrupted Trichomonas vaginalis |
| Isotype | IgG3 |
| Source/Host | Mouse |
| Species Reactivity | T. vaginalis |
| Clone | C986M |
| Affinity Constant | Not determined |
| Purification | 90% pure. Protein A chromatography |
| Conjugate | Unconjugated |
| Applications | Suitable for use in IFA and ELISA. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded. Recommended antibody pair for sandwich immunoassay: Capture Detection DMAB4429 DMAB4431 Suggested pair for testing (Capture - Detection): DMAB4429 - <u>DMAB4431</u> |
| Format | Purified, Liquid |
| Concentration | Lot specific (OD280nm, E0.1% = 1.3) |

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

© Creative Diagnostics All Rights Reserved

| Size | 1 mg |
|--------------|---|
| Buffer | 0.01M PBS, pH 7.2 |
| Preservative | 0.1% Sodium Azide |
| Storage | Short term (up to 2 weeks) store at 2-8°C. Long term, aliquot and store at -20°C. Avoid multiple freeze/thaw cycles |

BACKGROUND

| Introduction | Trichomonasvaginalisis an anaerobic, flagellated protozoan, a form of microorganism. Theparasitic microorganism is the causative agent of trichomoniasis, and is themost common pathogenic protozoan infection of humans in industrializedcountries. Infection rates between men and women are the same with womenshowing symptoms while infections in men are usually asymptomatic. Transmission takes place directly because the trophozoite does not have acyst. The WHO has estimated that 180 million cases of infection are acquiredannually worldwide. The estimates for North America alone are between 5 and 8million new infections each year, with an estimated rate of asymptomaticcases as high as 50%. Usually treatment consists of metronidazole andtinidazole. Trichomonasvaginalis is a protozoa which causes inflammation of the vaginal canal. It isconsidered to be a sexually transmitted disease. |
|--------------|---|
| Keywords | T vaginalis ; TV ; Trichomonas vaginalis; Eukarya; Metamonada; Parabasalia; Trichomonadida; Trichomonas; T. vaginalis |