



Anti-CIDEA (aa 1-190) polyclonal antibody (DPABH-03568)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Antigen	Description
,	-000:.p::0::

Apoptosis is related to many diseases and induced by a family of cell death receptors and their ligands. Cell death signals are transduced by death domain containing adapter molecules and members of the caspase family of proteases. These death signals finally cause the degradation of chromosomal DNA by activated DNase. DFF45/ICARD has been identified as inhibitor of caspase activated DNase DFF40/CAD. DFF45 related proteins CIDE A and CIDE B (for cell death inducing DFF like effector A and B) were recently identified. CIDE contains a new type of domain termed CIDE N, which has high homology with the regulatory domains of DFF45/ICAD and DFF40/CAD. Expression of CIDE A induces DNA fragmentation and activates apoptosis, which is inhibited by DFF45. CIDE A is a DFF45 inhibitable effector that promotes cell death and DNA fragmentation. CIDE A is expressed in many tissues.

Immunogen	Recombinant fragment corresponding to a region within amino acids 1-190 of Human CIDE A
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Purification	Immunogen affinity purified
Conjugate	Unconjugated
Applications	WB, IHC-P, ICC/IF
Format	Liquid
Size	100 μΙ
Buffer	pH: 7.00; Constituents: 0.75% Glycine, 1.21% Tris, 10% Glycerol

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Preservative	None
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

GENE INFORMATION

Gene Name	CIDEA cell death-inducing DFFA-like effector a [Homo sapiens]
Official Symbol	CIDEA
Synonyms	CIDEA; cell death-inducing DFFA-like effector a; CIDE-A; cell death activator CIDE-A;
Entrez Gene ID	1149
Protein Refseq	NP_001270.1
UniProt ID	<u>060543</u>
Function	protein homodimerization activity;