



Anti-NAT10 (aa 989-1000) polyclonal antibody (DPABH-03652)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Has protein acetyltransferase activity in vitro. Can acetylate both histones and microtubules. Histone acetylation may regulate transcription and mitotic chromosome de-condensation. Activates telomerase activity by stimulating the transcription of TERT, and may also regulate telomerase function by affecting the balance of telomerase subunit assembly, disassembly, and localization. Acetylates alpha-tubulin, which may affect microtubule stability and cell division.
Synthetic peptide: C-KSDKKRKLEAKQ, corresponding to internal sequence amino acids 989-1000 of Human NAT10
IgG
Goat
Human
Immunogen affinity purified
Unconjugated
ELISA, IHC-P
Liquid
50 μg
pH: 7.30; Constituents: 0.5% BSA, 99% Tris buffered saline
0.02% Sodium Azide
Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Email: info@creative-diagnostics.com

© Creative Diagnostics All Rights Reserved

GENE INFORMATION

Gene Name	NAT10 N-acetyltransferase 10 (GCN6-related) [Homo sapiens]
Official Symbol	NAT10
Synonyms	NAT10; N-acetyltransferase 10 (GCN5-related); ALP; NET43; N-acetyltransferase 10; N-acetyltransferase-like protein;
Entrez Gene ID	<u>55226</u>
Protein Refseq	NP 001137502.1
UniProt ID	Q9H0A0
Pathway	Ribosome biogenesis in eukaryotes; metapathway biotransformation;
Function	ATP binding; N-acetyltransferase activity; poly(A) RNA binding;