



Anti-SGK1 (C-terminal) polyclonal antibody (DPABH-15114)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Antigen	Description

Protein kinase that plays an important role in cellular stress response. Activates certain potassium, sodium, and chloride channels, suggesting an involvement in the regulation of processes such as cell survival, neuronal excitability and renal sodium excretion. Sustained high levels and activity may contribute to conditions such as hypertension and diabetic nephropathy. Mediates cell survival signals, phosphorylates and negatively regulates proapoptotic FOXO3A. Phosphorylates NEDD4L, which leads to its inactivation and to the subsequent activation of various channels and transporters such as ENaC, KCNA3/Kv1.3 or EAAT1. Isoform 2 exhibited a greater effect on cell plasma membrane expression of ENaC and Na(+) transport than isoform 1.

Immunogen	Synthetic peptide (Human) corresponding to the C-terminus of human SGK.
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Mouse, Rat, Human
Purification	Immunogen affinity purified
Conjugate	Unconjugated
Applications	WB, IP
Format	Liquid
Size	100 μg
Buffer	Constituents: 50% Glycerol, 1% BSA, PBS, pH 7.2
Preservative	0.02% Sodium Azide

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Email: info@creative-diagnostics.com

© Creative Diagnostics All Rights Reserved

cycles.

GENE INFORMATION

Official Symbol SGK1	
	ulated kinase 1; serum/glucocorticoid regulated kinase; SGK; Sgk1; serine/threonine protein kinase SGK; inase 1; SGK;
Entrez Gene ID 6446	
Protein Refseq NP 001137148	
UniProt ID <u>O00141</u>	
Chromosome Location 6q23	
	eabsorption; Class I PI3K signaling events; FoxO family r regulatory network; IL-6 Signaling Pathway; Insulin Pathway;
	gulator activity; chloride channel regulator activity; nucleotide lator activity; protein serine/threonine kinase activity; sodium