



# Anti-IGFBP6 polyclonal antibody [Biotin] (DPABY-697)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Antigen Description</b>	The superfamily of insulin-like growth factor (IGF) binding proteins include the six high-affinity IGF binding proteins (IGFBP) and at least four additional low-affinity binding proteins referred to as IGFBP related proteins (IGFBP-rP). All IGFBP superfamily members are cysteine-rich proteins with conserved cysteine residues, which are clustered in the amino- and carboxy-terminal thirds of the molecule. IGFBPs modulate the biological activities of IGF proteins. Some IGFBPs may also have intrinsic bioactivity that is independent of their ability to bind IGF proteins. Post-translational modifications of IGFBP, including glycosylation, phosphorylation and proteolysis, have been shown to modify the affinities of the binding proteins to IGF. ALS (Acid Labile Subunit) is a liver-derived protein that exists in a ternary complex with Insulin-like Growth Factor (IGF)-binding Protein-3 (IGFBP-3) or IGFBP-5, and either IGF-I or IGF-II. ALS increases the half-life of IGF/IGFBP complexes in circulation.
<b>Specificity</b>	Detects mouse IGFBP-6 in ELISAs and Western blots. In sandwich immunoassays, less than 0.1% cross-reactivity with recombinant human (rh) IGFBP-1, rhIGFBP-2, rhIGFBP-3, rhIGFBP-4, rhIGFBP-5, rhIGFBP-6 and rmlGF-I is observed.
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant mouse IGFBP-6 . Gly24-Gly238 Accession Number P47880
<b>Isotype</b>	IgG
<b>Source/Host</b>	Goat
<b>Species Reactivity</b>	Mouse
<b>Purification</b>	Antigen Affinity-purified
<b>Conjugate</b>	Biotin
<b>Applications</b>	Western Blot, ELISA Detection (Matched Pair)

<b>Procedure</b>	Matched Antibody Pairs
<b>Format</b>	Liquid
<b>Size</b>	50 µg
<b>Buffer</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.
<b>Preservative</b>	None
<b>Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">Igfbp6 insulin-like growth factor binding protein 6 [ Mus musculus (house mouse) ]</a>
<b>Official Symbol</b>	IGFBP6
<b>Synonyms</b>	IGFBP6; insulin-like growth factor binding protein 6; IGFBP-6; insulin-like growth factor-binding protein 6; IBP-6; IGF-binding protein 6; INSULIN-LIKE GROWTH FACTOR BINDING PROTEIN 6 PRECURSOR (IGFBP-6) (IBP-6) (IGF-BINDING PROTEIN 6);
<b>Entrez Gene ID</b>	<a href="#">16012</a>
<b>Protein Refseq</b>	<a href="#">NP_032370</a>
<b>UniProt ID</b>	<a href="#">P47880</a>
<b>Chromosome Location</b>	15 F3; 15
<b>Pathway</b>	Metabolism of proteins; Myometrial Relaxation and Contraction Pathways; Regulation of Insulin-like Growth Factor (IGF) transport and uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs);
<b>Function</b>	growth factor binding; insulin-like growth factor I binding; insulin-like growth factor II binding; insulin-like growth factor binding;