



Anti-c-Maf polyclonal antibody (DPATB-H81337)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

| Product Overview | Rabbit Anti-c-Maf Polyclonal Antibody |
|-----------------------------------|---|
| Target | c-Maf |
| Immunogen | Fusion protein. |
| Isotype | IgG |
| Source/Host | Rabbit |
| Species Reactivity | N/A |
| Purification | Whole antiserum |
| Conjugate | Unconjugated |
| Applications | WB, EMSA |
| Cellular Localization | Nucleus. |
| Post translation Modifications | Ubiquitinated, leading to its degradation by the proteasome. Ubiquitination is triggered by glucocorticoids. Phosphorylated by GSK3 and MAPK13 on serine and threonine residues (Probable). The phosphorylation status can serve to either stimulate or inhibit |
| Format | Liquid |
| Size | 200 μΙ |
| Buffer | Whole Serum |
| Preservative | None |
| | |

45-1 Ramsey Road, Shirley, NY 11967, USA

Email:info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

cycles.

BACKGROUND

Introduction Acts as a transcriptional activator or repressor. Involved in embryonic lens fiber cell

development. Recruits the transcriptional coactivators CREBBP and/or EP300 to crystallin

promoters leading to up-regulation of crystallin gene during lens fiber cell d

Keywords AS42 oncogene homolog; Avian musculoaponeurotic fibrosarcoma (MAF) protooncogene;

avian musculoaponeurotic fibrosarcoma (v maf); c Maf; c maf proto oncogene; cMaf; MAF;

MAF_HUMAN; MAF2; MGC71685; Proto oncogene c Maf; Proto-oncogene c-maf;

Transcription f

GENE INFORMATION

Entrez Gene ID 4094

UniProt ID <u>O75444</u>